

Bringing a Superposition $\alpha|11\rangle + \beta|00\rangle$ ($\alpha > \beta$) to a Bell State Through Quantum Zeno-like Measurements

J. G. Oliveira Jr., R. Rossi Jr., and M. C. Nemes

*Departamento de Física, Instituto de Ciências Exatas,
Universidade Federal de Minas Gerais, C.P. 702,
30161-970, Belo Horizonte, MG, Brazil*

Abstract

We show rather surprising features of Zeno-like measurements on two pairs of qubits system interacting pairwise. For the superposition state $\alpha|11\rangle + \beta|00\rangle$ ($\alpha > \beta$) and a finite set of measurements we find an entanglement enhancement. For $N \rightarrow \infty$ the initial state is preserved. Moreover we show that a single measurement after the “sudden death of entanglement” induces a new entangled state. For specific measurement times the entanglement will be maximum.

PACS numbers:

Keywords:

The possibility of freezing the evolution of a system by frequent measurements was firstly pointed out by von Neumann in 1932 [1], and formally presented by B. Misra and E. C. Sudarshan [2]. Although some experimental realizations of the effect have been reported [3, 4, 5, 6] it remained, for several years, the center of fervorous debates [7, 8, 9, 10, 11, 12]. Recently a conclusive experiment on an unstable quantum system was reported [13].

Interestingly enough Quantum Zeno-like measurements have become an important tool for practical applications in the growing field of quantum information theory. Error prevention [14, 15, 16, 17, 18], quantum state engineering [19], state preservation [20] and decoherence control [21, 22, 23] are some examples. Recently Zeno-like measurements have been used to protect an entangled state from its environment [24].

In the present contribution we show that Zeno-like measurements in the context of a double Jaynes-Cummings model [25] may not only preserve the maximum entanglement of a Bell state, but also, surprisingly, enhance the entanglement of the state $\alpha|11\rangle + \beta|00\rangle$ ($\alpha > \beta$), bringing it to a Bell state.

Many important features of entanglement dynamics have been extensively studied in such models, namely: relation between energy and entanglement [26], pairwise concurrence dynamics [27], entanglement invariant for this model [25] and sudden death of entanglement [28].

In this case (sudden death of entanglement) we were able to show, for this system, that a single measurement after the event may yield a new entangle state.

Let us consider four qubits system coupled $2(ab)$ by $2(AB)$ with identical coupling coefficient(g). The interaction is such that it couples a only to A , and b to B as in Jaynes-Cummings models quoted above.

Let us consider the initial state $|\phi_+\rangle_{ab}|0,0\rangle_{AB} = (\alpha_0|1,1\rangle_{ab} + \beta_0|0,0\rangle_{ab})|0,0\rangle_{AB}$, whose dynamical evolution is given as

$$\begin{aligned} |\phi_+\rangle_{ab}|0,0\rangle_{AB} \longrightarrow & (\alpha_0 a^2(t)|1,1\rangle_{ab} + \beta_0|0,0\rangle_{ab})|0,0\rangle_{AB} \\ & - \alpha_0 b(t) [ia(t)(|1,0\rangle_{ab}|0,1\rangle_{AB} + |0,1\rangle_{ab}|1,0\rangle_{AB}) + b(t)|0,0\rangle_{ab}|1,1\rangle_{AB}], \end{aligned} \quad (1)$$

where $a(t) = \cos(gt)$ and $b(t) = \sin(gt)$. Note that at $t = T = \frac{\pi}{2g}$ there is an entanglement swapping

$$|\phi_+\rangle_{ab}|0,0\rangle_{AB} \longrightarrow |0,0\rangle_{ab}|\phi_-\rangle_{AB}, \quad (2)$$

where $|0,0\rangle_{ab}|\phi_-\rangle_{AB} = |0,0\rangle_{ab}(\alpha_0|1,1\rangle_{AB} - \beta_0|0,0\rangle_{AB})$.

Let us introduce a probe system composed of N two level systems all prepared on $|0\rangle$ state, i.e. $(\bigotimes_{k=1}^N |0_M^{(k)}\rangle)$. This probe system interacts one at the time (at times $t_N = N\tau$) with subsystem AB . This interaction will discriminate, after each interaction, between the two possibilities: no excitation in AB and otherwise. We also assume that the probe system can be measured after every interaction with AB .

After a sequence of N interactions keeping only those with null result, i.e., the probe remains unexcited; the vector state will become

$$|\Phi_N(T)\rangle = \frac{1}{(|\alpha_0|^2 \cos^{4N}(g\tau) + |\beta_0|^2)^{1/2}} (\alpha_0 \cos^{2N}(g\tau) |1,1\rangle_{ab} + \beta_0 |0,0\rangle_{ab}) |0,0\rangle_{AB}, \quad (3)$$

where we fixed the total time of the evolutions as $T = \frac{\pi}{2g}$ (the entanglement swap time), and $\tau = \frac{T}{N}$.

It is apparent from this expression that the relative contribution of the state $|1,1\rangle_{ab}$ can be manipulated. In what follows we derive the conditions under which the entanglement enhancement of this system is possible.

For this purpose we calculate the concurrency [29] of the initial state

$$C_0 = \frac{2|\alpha_0||\beta_0|}{|\alpha_0|^2 + |\beta_0|^2} = 2|\alpha_0||\beta_0|, \quad (4)$$

and the one after N measurements

$$C_N = \frac{2|\alpha_0||\beta_0| \cos^{2N}(g\tau)}{|\alpha_0|^2 \cos^{4N}(g\tau) + |\beta_0|^2}, \quad (5)$$

Taking the limit $N \rightarrow \infty$ on (3) and (5), we can see the dynamics inhibition.

$$\lim_{N \rightarrow \infty} |\Phi_N(T)\rangle = |\phi_+\rangle_{ab}|0,0\rangle_{AB}, \quad (6)$$

$$\lim_{N \rightarrow \infty} C_N = C_0. \quad (7)$$

The vector state of the system after N measurements tends to its initial state as does the concurrence to its initial value when $N \rightarrow \infty$.

From equation (4) and the normalization of $|\phi_+\rangle_{ab}|0,0\rangle_{AB}$ ($|\alpha_0|^2 + |\beta_0|^2 = 1$), we may write:

$$|\alpha_0| = \begin{cases} \sqrt{\frac{1}{2} + \sqrt{\frac{1-C_0^2}{4}}}; & |\alpha_0| \geq |\beta_0| \\ \sqrt{\frac{1}{2} - \sqrt{\frac{1-C_0^2}{4}}}; & |\alpha_0| \leq |\beta_0| \end{cases} \quad (8)$$

Substituting (8) in (5), and after some algebra we may write the concurrence C_N as function of the initial concurrence and the number of measurements.

$$C_N^\pm = \begin{cases} C_N^+; & |\alpha_0| \geq |\beta_0| \\ C_N^-; & |\alpha_0| \leq |\beta_0| \end{cases} \quad (9)$$

where

$$C_N^\pm = \frac{2C_0 \cos^{2N}(g\tau)}{1 + \cos^{4N}(g\tau) \mp \sqrt{1 - C_0^2}[1 - \cos^{4N}(g\tau)]}. \quad (10)$$

The functions (C_N^+ and C_N^-) show a similar behavior as $N \rightarrow \infty$, both tend to the initial concurrence value. However, their behavior is quite different when N is finite, this is shown on Fig.1.

If only one measurement is performed at $\tau = \frac{\pi}{2g}$, no inhibition can be observed and the entanglement is completely transferred to the subsystem AB . Therefore, the concurrence of subsystem ab is null and both curves in Fig. 1 start from zero. As N increases C_N^- grows steadily tending to C_0 when $N \rightarrow \infty$. We conclude that for initial states with $|\alpha_0| \leq |\beta_0|$, the effect of entanglement freezing is the only one that can be induced.

On the other hand, the function C_N^+ shows an interesting effect besides the entanglement freezing. When the initial state respects the relation $|\alpha_0| > |\beta_0|$, the concurrence of the subsystem ab can exceed its initial value, and become very close to the maximum value of 1. This is best visualized in the coefficient $\alpha_0 \cos^{2N}(g\tau)$ (in (3)) which decreases as N increases, and when its value approaches β_0 the concurrence grows. If $\alpha_0 \cos^{2N}(g\tau) = \beta_0$ the concurrence is maximum $C_N = 1$.

It is also possible to bring the superposition $\alpha_0|11\rangle + \beta_0|00\rangle$ ($\alpha_0 > \beta_0$) to a Bell state with only one measurement. The procedure is simple, one just has to let the system evolve freely and perform a measurement on subsystem AB at the time $t = \frac{1}{g} \arccos(\sqrt{\beta_0/\alpha_0})$ (when




FIG. 1: Concurrence C_N^- (red) and C_N^+ (blue) as function of N for $\tau = \frac{\pi}{2gN}$.

$\alpha_0 a^2(t) = \beta_0$). Selecting the states with null result, one will have a Bell state prepared in subsystem ab .

The appearance of entanglement sudden death for the initial states $(\alpha|11\rangle + \beta|00\rangle)|0,0\rangle$ ($\alpha > \beta$), in the context of the double Jaynes-Cummings model is well known [28]. In order to explicitate this effect we give the expression for the concurrence (C_f) of the state evolving without external interference, as a function of the initial concurrence.

$$C_f^\pm(t) = \max(0, \Lambda^\pm(t)), \quad (11)$$

where

$$\Lambda^\pm(t) = \sqrt{1 \pm \sqrt{1 - C_0^2} \cos^2(gt)} \left(\sqrt{1 \mp \sqrt{1 - C_0^2}} - \sqrt{1 \pm \sqrt{1 - C_0^2} \sin^2(gt)} \right). \quad (12)$$

Notice that for times

$$t = \frac{1}{g} \arcsin \left(\frac{\sqrt[4]{1 - \sqrt{1 - C_0^2}}}{\sqrt[4]{1 + \sqrt{1 - C_0^2}}} \right), \quad (13)$$

which are prior to the swap time, there will be a sudden death of entanglement. Fig.2 illustrate this. However, in Fig.3 we show how a single measurement can induce an early entanglement resurrection.

The graphic on Fig.3 shows the concurrence (C_1^+) after a single measurement on AB with null result performed at times t , for different values of the initial concurrence of the states $(\alpha|11\rangle + \beta|00\rangle)|0,0\rangle$ ($\alpha > \beta$). Notice that (from Fig.2) the concurrence is zero after the sudden death time. In Fig.3, surprisingly enough, the set of vanishing final concurrences are “brought to life” by a single measurement performed not necessarily immediately after the time at which sudden death takes place.




FIG. 2: Concurrence without measurements (C_f) as function of gt and C_0 for initial states $(\alpha|11\rangle + \beta|00\rangle)|0,0\rangle$ ($\alpha > \beta$) .

The reason for this is that after the sudden death and before the entanglement swap time, there is no entanglement on ab , but the excitations are not completely transferred to AB . Then if one performs a measurement on AB , and gets a null result, a state with finite entanglement is prepared in ab . It is even possible to resurrect the entanglement to the maximum value ($C_1^+ = 1$), exceeding the initial one, if the measurement is performed at $t = \frac{1}{g} \arccos\left(\sqrt{\beta_0/\alpha_0}\right)$.

To summarize, we have shown that QZE is capable of enhancing the entanglement of the state $(\alpha|11\rangle + \beta|00\rangle)|0,0\rangle$ ($\alpha > \beta$). Entanglement freezing is possible for all the others Bell states, however not the enhancement. Also surprisingly is the fact that the sudden death, observed in other calculations [28], can be avoided by a single measurement.




FIG. 3: Concurrence with one measurement at time t , as function gt and C_0 for initial states $(\alpha|11\rangle + \beta|00\rangle)|0,0\rangle$ ($\alpha > \beta$) .

The authors acknowledge financial support by CNPq.

-
- [1] J. von Neumann, *Die Mathematische Grundlängen der Quantenmechanik*, Springer Verlag, Berlin, 1932, *Mathematical Foundations of Quantum Mechanics* Princeton University Press, 1955.
 - [2] B. Misra, E.C. G. Sudarshan, J. Math. Phys. **18**, 765 (1977).
 - [3] Wayne M. Itano, D. J. Heinzen, J. J. Bollinger, D. J. Wineland, Phys. Rev. A **41**, 2295 (1990).
 - [4] P. Kwiat, H. Weinfurter, T. Herzog, and A. Zeilinger, Phys. Rev. Lett. **74**, 4763 (1995).
 - [5] B. Nagels, L. J. F. Hermans, and P. L. Chapovsky, Phys. Rev. Lett. **79**, 3097 (1997).
 - [6] E. W. Streed, J. Mun, M. Boyd, G. K. Campbell, P. Medley, W. Ketterle and D. E. Pritchard, Phys. Rev. Lett. **97**, 260402 (2006).
 - [7] L. E. Ballentine Phys. Rev. A **43**, 5165 (1991).
 - [8] Wayne M. Itano, D. J. Heinzen, J. J. Bollinger, D. J. Wineland, Phys. Rev. A **43**, 5168 (1991).
 - [9] Saverio Pascazio, Mikio Namiki Phys. Rev. A **50**, 4582 (1994).
 - [10] P. Facchia, V. Gorinib, G. Marmoc, S. Pascazio, , a and E. C. G. Sudarshan, Physics Letters A **275**, 12 (2000).
 - [11] D. Home and M. A. B. Whitaker, Annals of Physics **258**, 237 (1997).
 - [12] P.E. Toschek and Ch. Wunderlich, Eur.Phys.J.D **14**, 387(2001).

- [13] M. C. Fischer, B. Gutiérrez-Medina, and M. G. Raizen, Phys. Rev. Lett. **87**, 040402 (2001).
- [14] Chui-Ping Yang and Shih-I Chu, Phys. Rev. A **66**, 034301 (2002).
- [15] L. Vaidman, L. Goldenberg, and S. Wiesner, Phys. Rev. A **54**, R1745 (1996).
- [16] Lu-Ming Duan and Guang-Can Guo, Phys. Rev. A **57**, 2399 (1998).
- [17] WonYoung Hwang, Hyukjae Lee, Doyeol (David) Ahn, and Sung Woo Hwang, Phys. Rev. Lett. **62**, 062305 (2000).
- [18] N. Erez, Y. Aharonov, B. Reznik, and L. Vaidman, Phys. Rev. A **69**, 062315 (2004).
- [19] M. Zhang and L. You, Phys. Rev. Lett. **91**, 230404 (2003).
- [20] D. Dhar, L. K. Grover, and S. M. Roy, Phys. Rev. Lett. **96**, 100405 (2006).
- [21] O. Hosten, M. T. Rakher, J. T. Barreiro, N. A. Peters and P. G. Kwiat, Nature **439**, 949 (2006).
- [22] J. D. Franson, B. C. Jacobs, and T. B. Pittman, Phys. Rev. A **70**, 062302 (2004).
- [23] P. Facchi, S. Tasaki, S. Pascazio, H. Nakazato, A. Tokuse and D. A. Lidar, Phys. Rev. A **71**, 022302 (2005).
- [24] S. Maniscalco, F. Francica, R. L. Zaffino, N. Lo Gullo and F. Plastina, Phys. Rev. Lett. **100**, 090503 (2008).
- [25] I. Sainz and G. Björk, Phys. Rev. A , **76**, 042313 (2007).
- [26] D. Cavalcanti, J. G. Oliveira Jr, J. G. P. de Faria, M. O. T. Cunha and M. F. Santos, Phys. Rev. A , **74**, 042328 (2006).
- [27] M. Yönaç, T. Yu and J. H. Eberly, J. Phys. B, **40**, S45 (2007).
- [28] M. Yönaç, T. Yu and J. H. Eberly, J. Phys. B, **39**, S621 (2006).
- [29] S. Hill and W. K. Wootters, Phys. Rev. Lett. , **78**, 5022 (1997).